Tee Time®

0-0-30

DIRECTIONS FOR USE

Apply only as directed during the growing season.

Application Rates:

1.23 - 2.03 kg/100 m² (2.50 - 4.15 lbs/1,000 sq. ft.)

Apply at the rate for the amount of Potassium desired.

Use Rate kg/100 m ²	K ₂ O kg/100 m ²	Use Rate lbs/1,000 sq. ft.	K ₂ O lbs/1,000 sq. ft.	20 kg bag treats	20 kg bag treats	
1.23	0.37	2.50	0.75	1,620 m ²	17,600 sq. ft.	
1.63	0.49	3.33	1.00	1,220 m ²	13,200 sq. ft.	
2.03	0.61	4.15	1.25	980 m ²	10,600 sq. ft.	

For all applications:

- Apply to dry foliage.
- Water in after application with a minimum of 5.1 mm (0.2 inches) of irrigation to help move particles into the turf canopy.
- Do not collect clippings for at least one mowing following application to avoid particle pick up.
- If temperatures exceed 29° C (85° F) or if application to damp or wet turf is unavoidable, irrigate immediately after application.
- Apply at low rates if temperatures exceed 29° C (85° F).

Notice: This product contains the micronutrient iron for extra greening and to help prevent iron deficiency. Iron may stain concrete and other surfaces. Misapplied product to these surfaces should be removed promptly by sweeping or blowing. **DO NOT WASH OFF WITH WATER**.

Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn and sweep any product that lands on the driveway, sidewalk or street, back onto your lawn.

GUARANTEED ANALYSIS

	30.00%		
Sulfur (S)	10.20%		
Copper (Cu)	0.10%		
0.02% Water Soluble Copper (Cu)	*****		
Iron (Fe)	3.00%		
0.03% Water Soluble Iron (Fe)	0.10%		
Manganese (Mn)			
0.03% Water Soluble Manganese (Mn)			
Zinc (Zn)	0.10%		
0.03% Water Soluble Zinc (Zn)			

Constituent Materials: potassium sulfate, cupric oxide, cupric sulfate, ferric oxide, ferrous sulfate, manganous oxide, manganese sulfate, zinc oxide and zinc sulfate.

Storage and Use:

Store in a clean, dry place.
Reseal opened bag by folding top down and securing.
Use only clean, dry spreaders.
Do not leave unused product in spreaders.

Notice: Warranty of this material, either expressed or implied, is limited to a guarantee of the composition as shown on the label in as much as uses are beyond the seller's control. For the same reason, Seller is not liable for any injury to living things, crops, soils or materials which may result from the use or storage of this material. Do not store outside.

Product of USA

The Andersons logo and Tee Time are registered trademarks of The Andersons, Inc.; MUtech is a trademark of The Andersons, Inc.

Suggested Spreader Settings The suggested spreader settings are not intended to replace calibration. Please calibrate your spreader before applying product.									
SPREADER	GROUND SPEED	WIDTH OF COVERAGE	1.23 kg/100 m ² (2.50 lbs/1,000 sq. ft.)	1.63 kg/100 m ² (3.33 lbs/1,000 sq. ft.)	2.03 kg/100 m ² (4.15 lbs/1,000 sq. ft.)				
Model 2000 (cone 9), 2000 SR (cone 9)	4.8 km/hr (3 mph)	2.7 m (9 feet)	I 1/2	J 1/2	K 1/4				
SSD Drop	4.8 km/hr (3 mph)	overlap wheels	3 1/4	3 3/4	4 1/4				
R-8, R-8A (cone 9)	4.8 km/hr (3 mph)	2.7 m (9 feet)	H 1/4	ı	J				
Lely Models WTR, WFR, HR, 1250 (PTO at 450 rpm) (Pattern III-A)	7.2 km/hr (4.5 mph)	8.5 m (28 feet)	3 1/4	3 3/4	4 1/4				
Vicon (03 Series)	7.2 km/hr (4.5 mph)	7.3 m (24 feet)	13	16	19				

Manufactured by: The Andersons, Inc. P. O. Box 119 Maumee, OH 43537 USA www.andersonsplantnutrient.com





Net Weight 20 kg (44.1 lbs.)
APPCSOPM4

C18

This section of label reserved for French translation